

Physical Principles in Biology

Biology 3550

Spring 2024

Lecture 29

Enzymatic Coupling and

Introduction to Water

Wednesday, 27 March

©David P. Goldenberg


University of Utah

goldenberg@biology.utah.edu

Announcements


- Problem Set 4:
 - Due Monday, 1 April at 11:59 PM
 - Submit pdf file on Gradescope
- No office hours on Thursday, 28 March
- Review Session:
 - 5:15 PM, Thursday, 28 March
 - HEB 2002
- Quiz 4:
 - Friday, 29 March
 - 25 min, second half of class
 - Will cover thermodynamics
 - 50 min
- No class on Monday, 8 April

ATP Hydrolysis




- $\Delta G^\circ = -30 \text{ kJ/mol}$, $K_{\text{eq}} \approx 2 \times 10^5 \text{ M}$
- Why is the reaction so favorable?
 - High density of negative phosphate charges is reduced.
 - More resonance stabilization in ADP and P_i .
 - More favorable interaction with water by ADP and P_i .

Kinetics of ATP Hydrolysis




- Half-time is ≈ 20 days at neutral pH and 60°C .
- Transition state is a high energy state with equal probability of breaking down in either direction.
- Reaction rate is proportional to probability of forming the transition state.
- See the lobby of the Henry Eyring Building!

Catalysis of ATP Hydrolysis




- Enzymes catalyze chemical reactions by lowering transition-state energy.
- Enzymes create micro-environments that favor forming the transition state.
- Simply catalyzing ATP hydrolysis is not useful!

Enzymatic Coupling




- Enzyme mechanistically couples reactions.
- $\text{ATP} \longrightarrow \text{ADP} + \text{P}_i$ can't occur without $\text{A} \longrightarrow \text{B}$
- $\text{A} \longrightarrow \text{B}$ can't occur without $\text{ATP} \longrightarrow \text{ADP} + \text{P}_i$
- Coupled "reaction" can be physical motion or transport across membranes.


Enzymatic Coupling: Creatine Kinase



■ Coupled reactions





Hydrolysis of Creatine Phosphate



$\Delta G^\circ = -43 \text{ kJ/mol}$; more favorable than ATP hydrolysis.

Clicker Question #1


In which direction is the reaction favorable under standard-state conditions?



C) Neither direction is more favorable than the other.


Enzymatic Coupling: Creatine Kinase

■ Hydrolysis of creatine phosphate:



$\Delta G^\circ = -43 \text{ kJ/mol}$; more favorable than ATP hydrolysis.


■ Coupled reactions



Phosphorylation of ADP by creatine phosphate is favored.

(at standard-state concentrations!)

Creatine Kinase in Muscle Cells



- Typical concentrations in resting muscle cells:

4 mM ATP

0.013 mM ADP

25 mM creatine phosphate

13 mM creatine


- Calculate ΔG for these concentrations:

$$\begin{aligned}\Delta G &= \Delta G^\circ + RT \ln \frac{[\text{ATP}][\text{C}]}{[\text{ADP}][\text{CP}]} \\ &= -13 \text{ kJ/mol} + RT \ln \frac{0.004 \text{ M} \cdot 0.013 \text{ M}}{1.3 \times 10^{-5} \text{ M} \cdot 0.025 \text{ M}}\end{aligned}$$

$$\Delta G \approx 0$$

- Because enzyme keeps reaction at equilibrium!

Creatine Kinase in Muscle Cells



- If 1 mM ATP is suddenly converted to ADP:

4 mM ATP \rightarrow 3 mM ATP

0.013 mM ADP \rightarrow 1 mM ADP

25 mM creatine phosphate

13 mM creatine

- Calculate ΔG for these concentrations:


$$\Delta G = \Delta G^\circ + RT \ln \frac{[\text{ATP}][\text{C}]}{[\text{ADP}][\text{CP}]}$$

$$= -13 \text{ kJ/mol} + RT \ln \frac{0.003 \text{ M} \cdot 0.013 \text{ M}}{0.001 \text{ M} \cdot 0.025 \text{ M}}$$

$$\approx -12 \text{ kJ/mol}$$

- Creatine phosphate provides emergency reserve of free energy.

Warning!




Direction Change

Water: What Makes it Special?

- Unusually high boiling temperature.
 - Boiling temperature is point where vapor pressure reaches atmospheric pressure.
 - Generally, boiling temperature reflects strength of interactions between molecules.
 - Generally, boiling temperature increases with size of molecules, because larger molecules have larger surfaces for interaction.
 - Boiling temperature of water is very high for its size.
- Melting temperature of solid (ice) is also relatively high for size.
- Does not mix well with many other liquids, especially hydrocarbons.

Water Molecules are Polar


- Chemical bonds represent “sharing” of electrons between atoms.
- In some bonds, sharing is quite even: C-C, O-O, C-H
- Some elements are “greedy” for electrons: Electronegative elements: Oxygen and Nitrogen
- Some elements are “generous” with electrons: Electropositive elements: H, metals
- Water is particularly polar:



Oxygen has partial negative charge, and hydrogens have partial positive charges.


Hydrogen Bonds

- Form between a hydrogen atom covalently bonded to an electronegative atom and a second electronegative atom.



- Electronegative atoms are usually nitrogen or oxygen in biological molecules.
- Largely accounted for by electrostatic interaction between partial positive and negative charges, but there is probably also a small degree of covalent character to hydrogen bonds.
- Significant variability in geometry and strength of hydrogen bonds.

Hydrogen Bonds Between Water Molecules



- Each water molecule can act as a donor for two hydrogen bonds and an acceptor for two hydrogen bonds.
- In ice, each water molecule is hydrogen bonded to four others. Same geometry as carbon atoms in a diamond.
- What about in liquid water?

Clicker Question #2

How many hydrogen bonds does a water molecule form, on average, in liquid water at room temperature?

A) 0

B) 1


C) 2

D) 3

E) 4

All answers count for now.

Hydrogen Bonds in Liquid Water



- On average, each water molecule forms 3 hydrogen bonds at any instant.
- Explains high boiling point of water.
- Hydrogen bonds break and form constantly, giving water liquid properties.

Picture from simulation by
Prof. Valeria Molinero
U of U Chemistry Department.