

Maxima workbook for Principles of NMR Spectroscopy

Chapter 12: More quantum mechanics: Time and energy

1 Introduction

This wxMaxima workbook is an electronic supplement to to the book Principles of NMR Spectroscopy: An Illustrated Guide, David P. Goldenberg, University Science Books, (c) 2016
This and related files are available for download through links at: <http://uscibooks.com/goldenberg.htm>
wxMaxima is a graphical user interface to the computer algebra system (CAS) Maxima. General information about Maxima and wxMaxima, along with free versions of the programs, can be found at: <http://maxima.sourceforge.net/> and <http://andrejv.github.io/wxmaxima/>
Before attempting to use this workbook, users are strongly encouraged to read and experiment with the introductory workbook, gettingStarted.wmx, and the workbook for Chapter 11, chapter11.wmx
This software is distributed under the conditions of the BSD license and without any guarantees or warranties. (c) 2016 by David P. Goldenberg
Please send comments, including bug reports, to this address:
David P. Goldeberg
Department of Biology
University of Utah
257 South 1400 East
Salt Lake City, UT 84112-0840
goldenberg@biology.utah.edu

This chapter covers the general description of the change of a wavefunction with time and the specific cases of a spin-1/2 particle in a stationary field and during a pulse.
The worksheet uses the definitions found in 1spinLib.mac

```
(%i1) load("1spinLib.mac")$
```

The list of functions defined in this macro file (and the packages it uses, can be generated with the functions command.

```
(%i2) functions;
```

```
(%o2) [innerproduct (x, y) , unitvector (x) , columnvector (x) , gramschmidt (x, [myinnerproduct]) , eigenvalues (mat) , eigenvectors (mat) , sub
```

2 12.2 Spin-1/2 particles in a magnetic field

Page 348

3 12.2.1 Time dependence of the wave function

Here, we express the Hamiltonian operator as a matrix. The dependence on the external field strength and gyromagnetic ratio are expressed through the Larmor frequency, $\nu = -\gamma B / (2\pi)$

```
(%i3) H(nu);
```

```
(%o3)  
$$\begin{pmatrix} \frac{h\nu}{2} & 0 \\ 0 & -\frac{h\nu}{2} \end{pmatrix}$$

```

The general form of a wavefunction, or ket, for a spin-1/2 particle is written as a column vector:

```
(%i4) k_psi;
```

```
(%o4)  
$$\begin{pmatrix} ca \\ cb \end{pmatrix}$$

```

The complex conjugate, bra, of the wavefunction is written as a row vector:

```
(%i5) b_psi;
```

$$(\%o5) \quad \left(\overline{ca} \quad \overline{cb} \right)$$

The operation of the Hamiltonian on the wavefunction

$$(\%i6) \text{ H}(\text{nu}).\text{k_psi};$$

$$(\%o6) \quad \left(\begin{array}{c} \frac{ca \cdot h \cdot \nu}{2} \\ -\frac{cb \cdot h \cdot \nu}{2} \end{array} \right)$$

The eigenvalues and eigenvectors of the Hamiltonian

$$(\%i7) \text{ uniteigenvectors}(\text{H}(\text{nu}));$$

$$(\%o7) \quad [[[-\frac{h \cdot \nu}{2}, \frac{h \cdot \nu}{2}], [1, 1]], [[[0, 1]], [[1, 0]]]]$$

The eignenvectors of H are the same as those of Iz and the eigenvalues are -h*nu/2 and h*nu/2. For a nucleus with a positive gyromagnetic ratio, the frequency is negative and energy of the lbeta> wavefunction (represented as a vector in Maxima list form as [0,1]) has a positive energy.

The time dependence of a wavefunction is calculated using the unitary matrix Uh, which includes terms for both the time period, t, and the Larmor frequency nu

$$(\%i8) \text{ Uh}(\text{t}, \text{nu});$$

$$(\%o8) \quad \left(\begin{array}{cc} e^{-i \cdot \pi \cdot \nu \cdot t} & 0 \\ 0 & e^{i \cdot \pi \cdot \nu \cdot t} \end{array} \right)$$

The product of this matrix and the vector form of a wavefunction gives the wavefunction after time, t

$$(\%i9) \text{ Uh}(\text{t}, \text{nu}).\text{k_psi};$$

$$(\%o9) \quad \left(\begin{array}{c} ca \cdot e^{-i \cdot \pi \cdot \nu \cdot t} \\ cb \cdot e^{i \cdot \pi \cdot \nu \cdot t} \end{array} \right)$$

We use as the starting wavefunction one of the eigenfunctions of lx

$$(\%i10) \text{ k_0:ket}(1/\text{sqrt}(2), 1/\text{sqrt}(2));$$

$$(\%o10) \quad \left(\begin{array}{c} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{array} \right)$$

The complex conjugate of the wavefunction (i.e. the bra) is

$$(\%i11) \text{ b_0:bra}(\text{k_0});$$

$$(\%o11) \quad \left(\begin{array}{cc} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{array} \right)$$

After time t, the wavefunction is

$$(\%i12) \text{ Uh}(\text{t}, \text{nu}).\text{k_0};$$

$$(\%o12) \quad \left(\begin{array}{c} \frac{e^{-i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \\ \frac{e^{i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \end{array} \right)$$

An aside on complex numbers. If we define a complex number as:

$$(\%i13) \text{ declare}(\text{c}, \text{complex});$$

$$(\%o13) \quad done$$

$$(\%i14) \text{ c:a + \%i*b};$$

(%o14) $i \cdot b + a$

(%i15) `c;`

(%o15) $i \cdot b + a$

The exponential form of the complex number can be found using the polarform command, found as the Convert to Polar Form command in the Simplify>>Complex menu of wxMaxima

(%i16) `cp:polarform(c);`

(%o16) $\sqrt{b^2 + a^2} \cdot e^{i \cdot \text{atan2}(b,a)}$

(%i17) `cp;`

(%o17) $\sqrt{b^2 + a^2} \cdot e^{i \cdot \text{atan2}(b,a)}$

The term sqrt(b^2+a^2) represents the modulus, m, and atan2(b,a) is the angle theta. The function atan2 calculates the arc tangent (inverse tangent) of points defined by their x and y coordinates.

(%i18) `? atan2;`

-- Function: atan2 (<y>, <x>)
- yields the value of 'atan(<y>/<x>)' in the interval '-\%pi' to
\%pi'.

(%o18) true

Complex numbers in the exponential form can also be converted to the rectangular form (Convert to Rectangular Form in the Simplify>>Complex menu of wxMaxima).

(%i19) `rectform(cp);`

(%o19) $i \cdot b + a$

4 12.2.2 Time dependence of the z-magnetization

Page 351.

We will call the evolving wavefunction, calculated above, k_t

(%i20) `k_t:Uh(t,nu).k_0;`

(%o20) $\begin{pmatrix} \frac{e^{-i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \\ \frac{e^{i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \end{pmatrix}$

The corresponding bra is b_t and can be formed using the bra function in 1spinLib.mac

(%i21) `b_t:bra(k_t);`

(%o21) $\begin{pmatrix} \frac{e^{i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} & \frac{e^{-i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \end{pmatrix}$

The z-magnetization is calculated by multiplying the ket by the Iz operator matrix, and then multiplying the product by the bra

(%i22) `Iz;`

(%o22) $\begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix}$

(%i23) `b_t.Iz.k_t;`

(%o23) 0

For the more general case, we can use k_psi and b_psi defined in 1spinLib.mac

```
(%i24) k_psi;
```

```
(%o24)      ⎛  ca ⎞
            ⎜  cb ⎟
```

```
(%i25) b_psi;
```

```
(%o25)      ⎛  cā   cb̄ ⎞
```

The over scores indicate the complex conjugates of ca and cb
The initial z-magnetization is:

```
(%i26) b_psi.Iz.k_psi;
```

```
(%o26)      ca · cā   cb · cb̄
            2       2
```

The time-dependent wavefunction and its complex conjugate are then:

```
(%i27) k_psit:Uh(t,nu).k_psi;
```

```
(%o27)      ⎛  ca · e-i·π·ν·t ⎞
            ⎜  cb · ei·π·ν·t ⎟
```

```
(%i28) b_psit:bra(k_psit);
```

```
(%o28)      ⎛  cā · ei·π·ν·t   cb̄ · e-i·π·ν·t ⎞
```

The time-dependent z-magnetization is

```
(%i29) b_psit.Iz.k_psit;
```

```
(%o29)      ca · cā   cb · cb̄
            2       2
```

Thus, the z-magnetization does not change with time, in the absence of relaxation or other perturbations.

The magnetization components can also be calculated using the meanPsi function in 1spinLib.mac.

```
(%i30) fundef(meanPsi);
```

```
(%o30)      meanPsi(op,ψ):= trigreduce (trigsimp (demoivre (ψ̄.op.ψ)))
```

The functions trigreduce, trigsimp and demoivre are included in this function definition to simplify the results and express them in trigonometric form.

```
(%i31) meanPsi(Iz,k_t);
```

```
(%o31)      0
```

```
(%i32) meanPsi(Iz,k_psi);
```

```
(%o32)      - cb · cb̄ - ca · cā
            2
```

```
(%i33) meanPsi(Iz,k_psit);
```

```
(%o33)      - cb · cb̄ - ca · cā
            2
```

5 12.2.3 Time dependence of the x- and y-magnetization components

The Ix operator matrix is:

(%i34) $I_x;$

(%o34)
$$\begin{pmatrix} 0 & \frac{1}{2} \\ \frac{1}{2} & 0 \end{pmatrix}$$

Using the previous definitions of k_t and b_t

(%i35) $k_t;$

(%o35)
$$\begin{pmatrix} \frac{e^{-i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \\ \frac{e^{i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \end{pmatrix}$$

(%i36) $b_t;$

(%o36)
$$\begin{pmatrix} \frac{e^{i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} & \frac{e^{-i \cdot \pi \cdot \nu \cdot t}}{\sqrt{2}} \end{pmatrix}$$

The initial x-magnetization is:

(%i37) $b_0.I_x.k_0;$

(%o37)
$$\frac{1}{2}$$

The time-dependent x-magnetization is:

(%i38) $b_t.I_x.k_t;$

(%o38)
$$\frac{e^{2 \cdot i \cdot \pi \cdot \nu \cdot t}}{4} + \frac{e^{-2 \cdot i \cdot \pi \cdot \nu \cdot t}}{4}$$

This can be simplified and converted to a trigonometric form

(%i39) $rectform(%);$

(%o39)
$$\frac{\cos (2 \cdot \pi \cdot \nu \cdot t)}{2}$$

The conversion is done automatically by the meanPsi function

(%i40) $meanPsi(I_x,k_t);$

(%o40)
$$\frac{\cos (2 \cdot \pi \cdot \nu \cdot t)}{2}$$

For the y-magnetization

(%i41) $I_y;$

(%o41)
$$\begin{pmatrix} 0 & -\frac{i}{2} \\ \frac{i}{2} & 0 \end{pmatrix}$$

The initial y=magnetization

(%i42) $meanPsi(I_y,k_0);$

(%o42)
$$0$$

The time-dependent y-magnetization

(%i43) b_t.Iy.k_t;

(%o43)
$$\frac{i \cdot e^{-2 \cdot i \cdot \pi \cdot \nu \cdot t}}{4} - \frac{i \cdot e^{2 \cdot i \cdot \pi \cdot \nu \cdot t}}{4}$$

(%i44) meanPsi(Iy,k_t);

(%o44)
$$\frac{\sin(2 \cdot \pi \cdot \nu \cdot t)}{2}$$

The 1spinLib.mac library also includes a function, allMagPsi, to calculate the x, y and z magnetization components in one command

(%i45) allMagPsi(k_psi);

$$\langle I_x \rangle = \frac{\overline{ca} \cdot cb + ca \cdot \overline{cb}}{2} \quad \langle I_y \rangle = \frac{i \cdot ca \cdot \overline{cb} - i \cdot \overline{ca} \cdot cb}{2} \quad \langle I_z \rangle = -\frac{cb \cdot \overline{cb} - ca \cdot \overline{ca}}{2}$$

(%o45)

(%i46) allMagPsi(k_0);

$$\langle I_x \rangle = \frac{1}{2} \quad \langle I_y \rangle = 0 \quad \langle I_z \rangle = 0$$

(%o46)

(%i47) allMagPsi(k_t);

$$\langle I_x \rangle = \frac{\cos(2 \cdot \pi \cdot \nu \cdot t)}{2} \quad \langle I_y \rangle = \frac{\sin(2 \cdot \pi \cdot \nu \cdot t)}{2} \quad \langle I_z \rangle = 0$$

(%o47)

(%i48) allMagPsi(k_psit);

$$\langle I_x \rangle = -\frac{-\overline{ca} \cdot cb \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) - ca \cdot \overline{cb} \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) - i \cdot \overline{ca} \cdot cb \cdot \sin(2 \cdot \pi \cdot \nu \cdot t) + i \cdot ca \cdot \overline{cb} \cdot \sin(2 \cdot \pi \cdot \nu \cdot t)}{2} \quad \langle I_y \rangle = \frac{-i \cdot \overline{ca} \cdot cb \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) + i \cdot ca \cdot \overline{cb} \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) - \overline{ca} \cdot cb \cdot \sin(2 \cdot \pi \cdot \nu \cdot t) + ca \cdot \overline{cb} \cdot \sin(2 \cdot \pi \cdot \nu \cdot t)}{2}$$

(%o48)

The simplification rules included in the allMagPsi function don't always give the simplest results, but further manipulations can be applied to the results. For instance for the x-magnetization from k_psit

(%i49) meanPsi(Ix,k_psit);

(%o49)
$$-\frac{-\overline{ca} \cdot cb \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) - ca \cdot \overline{cb} \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) - i \cdot \overline{ca} \cdot cb \cdot \sin(2 \cdot \pi \cdot \nu \cdot t) + i \cdot ca \cdot \overline{cb} \cdot \sin(2 \cdot \pi \cdot \nu \cdot t)}{2}$$

(%i50) ratsimp(%);

(%o50)
$$-\frac{\left(-ca \cdot \overline{cb} - \overline{ca} \cdot cb\right) \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) + \left(i \cdot ca \cdot \overline{cb} - i \cdot \overline{ca} \cdot cb\right) \cdot \sin(2 \cdot \pi \cdot \nu \cdot t)}{2}$$

This still looks pretty messy, but notice that the initial lx and ly components are:

(%i51) meanPsi(Ix,k_psi);

(%o51)
$$\frac{\overline{ca} \cdot cb + ca \cdot \overline{cb}}{2}$$

(%i52) meanPsi(Iy,k_psi);

(%o52)
$$\frac{i \cdot ca \cdot \overline{cb} - i \cdot \overline{ca} \cdot cb}{2}$$

If we call the initial values lx0 and ly0, we can substitute these values into the expression for the time-dependent lx magnetization

First, let's assign the expression to a variable.

```
(%i53) IxTime:ratsimp(meanPsi(Ix,k_psi));
```

$$(\%o53) \quad - \frac{\left(-ca \cdot \overline{cb} - \overline{ca} \cdot cb\right) \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) + \left(i \cdot ca \cdot \overline{cb} - i \cdot \overline{ca} \cdot cb\right) \cdot \sin(2 \cdot \pi \cdot \nu \cdot t)}{2}$$

Then we can use the substitution command, which is available as Simplify>>Substitute in wxMaxima. We do this in two steps.

```
(%i54) subst(2*Iy0, (%i*ca*conjugate(cb)-%i*conjugate(ca)*cb), IxTime);
```

$$(\%o54) \quad - \frac{\left(-ca \cdot \overline{cb} - \overline{ca} \cdot cb\right) \cdot \cos(2 \cdot \pi \cdot \nu \cdot t) + 2 \cdot Iy0 \cdot \sin(2 \cdot \pi \cdot \nu \cdot t)}{2}$$

```
(%i55) subst(-2*Ix0, (-ca*conjugate(cb)-conjugate(ca)*cb), %);
```

$$(\%o55) \quad - \frac{2 \cdot Iy0 \cdot \sin(2 \cdot \pi \cdot \nu \cdot t) - 2 \cdot Ix0 \cdot \cos(2 \cdot \pi \cdot \nu \cdot t)}{2}$$

Substitutions are rather tricky, and the form used in the substitution often has to match exactly the form that appears in the original expression.

6 12.3 When the Hamiltonian changes with time: The effect of radiation on the wavefunction

Page 356.

6.1 An x-pulse

Page 357.

The effects of pulses on the wavefunction are expressed in matrix form by rotation matrices containing the variable a, for the angle of rotation. For an x-pulse, the rotation matrix is (from 1spinLib.mac):

```
(%i56) Rx(a);
```

$$(\%o56) \quad \begin{pmatrix} \cos\left(\frac{a}{2}\right) & -i \cdot \sin\left(\frac{a}{2}\right) \\ -i \cdot \sin\left(\frac{a}{2}\right) & \cos\left(\frac{a}{2}\right) \end{pmatrix}$$

To obtain the new wavefunction, the existing wavefunction is multiplied by the rotation matrix. (Eq 12.6, page 359)

```
(%i57) Rx(a).k_psi;
```

$$(\%o57) \quad \begin{pmatrix} \cos\left(\frac{a}{2}\right) \cdot ca - i \cdot \sin\left(\frac{a}{2}\right) \cdot cb \\ \cos\left(\frac{a}{2}\right) \cdot cb - i \cdot \sin\left(\frac{a}{2}\right) \cdot ca \end{pmatrix}$$

1spinLib.mac also contains a function to calculate this result

```
(%i58) psiPulseX(k_psi,a);
```

$$(\%o58) \quad \begin{pmatrix} \cos\left(\frac{a}{2}\right) \cdot ca - i \cdot \sin\left(\frac{a}{2}\right) \cdot cb \\ \cos\left(\frac{a}{2}\right) \cdot cb - i \cdot \sin\left(\frac{a}{2}\right) \cdot ca \end{pmatrix}$$

Additional functions are provided for the specific cases of pi and pi/2 pulses along the x-axis

```
(%i59) psiPiX(k_psi);
```

$$(\%o59) \quad \begin{pmatrix} -i \cdot cb \\ -i \cdot ca \end{pmatrix}$$

(%i60) psiPi2X(k_psi);

(%o60)
$$\begin{pmatrix} \frac{ca}{\sqrt{2}} - \frac{i \cdot cb}{\sqrt{2}} \\ \frac{cb}{\sqrt{2}} - \frac{i \cdot ca}{\sqrt{2}} \end{pmatrix}$$

For the case of a (pi/2)x pulse applied to lalpha>

(%i61) psiPi2X(k_a);

(%o61)
$$\begin{pmatrix} \frac{1}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} \end{pmatrix}$$

(%i62) allMagPsi(%);

$$\langle I_x \rangle = 0 \quad \langle I_y \rangle = -\frac{1}{2} \quad \langle I_z \rangle = 0$$

(%o62)

6.2 y-pulses

Page 360.

The y-rotation matrix is:

(%i63) Ry(a);

(%o63)
$$\begin{pmatrix} \cos\left(\frac{a}{2}\right) & -\sin\left(\frac{a}{2}\right) \\ \sin\left(\frac{a}{2}\right) & \cos\left(\frac{a}{2}\right) \end{pmatrix}$$

(%i64) Ry(a).k_psi;

(%o64)
$$\begin{pmatrix} \cos\left(\frac{a}{2}\right) \cdot ca - \sin\left(\frac{a}{2}\right) \cdot cb \\ \cos\left(\frac{a}{2}\right) \cdot cb + \sin\left(\frac{a}{2}\right) \cdot ca \end{pmatrix}$$

y-pulse functions

(%i65) psiPulseY(k_psi,a);

(%o65)
$$\begin{pmatrix} \cos\left(\frac{a}{2}\right) \cdot ca - \sin\left(\frac{a}{2}\right) \cdot cb \\ \cos\left(\frac{a}{2}\right) \cdot cb + \sin\left(\frac{a}{2}\right) \cdot ca \end{pmatrix}$$

(%i66) psiPi2Y(k_psi);

(%o66)
$$\begin{pmatrix} \frac{ca}{\sqrt{2}} - \frac{cb}{\sqrt{2}} \\ \frac{cb}{\sqrt{2}} + \frac{ca}{\sqrt{2}} \end{pmatrix}$$

(%i67) psiPiY(k_psi);

(%o67)
$$\begin{pmatrix} -cb \\ ca \end{pmatrix}$$

(%i68) psiPulseX(k_a,a);

(%o68)
$$\begin{pmatrix} \cos\left(\frac{a}{2}\right) \\ -i \cdot \sin\left(\frac{a}{2}\right) \end{pmatrix}$$

A (pi/2)y pulse applied to lalpha>


```
(%i69) psiPi2Y(k_a);
```

(%o69)
$$\begin{pmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{pmatrix}$$

```
(%i70) allMagPsi(%);
```

$$\langle I_x \rangle = \frac{1}{2} \langle I_y \rangle = 0 \quad \langle I_z \rangle = 0$$

(%o70)

To look at the effect of an x-pulse after aligning the magnetization with the y'-axis, first define the wavefunction following a pi/2 x-pulse to |alpha>

```
(%i71) k_y:psiPi2X(k_a);
```

(%o71)
$$\begin{pmatrix} \frac{1}{\sqrt{2}} \\ -\frac{i}{\sqrt{2}} \end{pmatrix}$$

```
(%i72) allMagPsi(k_y);
```

$$\langle I_x \rangle = 0 \quad \langle I_y \rangle = -\frac{1}{2} \quad \langle I_z \rangle = 0$$

(%o72)

then apply a pi/2 y-pulse

```
(%i73) psiPi2Y(k_y);
```

(%o73)
$$\begin{pmatrix} \frac{i}{2} + \frac{1}{2} \\ \frac{1}{2} - \frac{i}{2} \end{pmatrix}$$

```
(%i74) allMagPsi(%);
```

$$\langle I_x \rangle = 0 \quad \langle I_y \rangle = -\frac{1}{2} \quad \langle I_z \rangle = 0$$

(%o74)

As expected, the second pulse has no effect.

6.3 A 2*Pi-pulse

Page 362.

A 2*pi pulse applied to |alpha>

```
(%i75) psiPulseX(k_a,2*%pi);
```

(%o75)
$$\begin{pmatrix} -1 \\ 0 \end{pmatrix}$$

The sign of the wavefunction has been reversed, but the observable magnetization components are unchanged.

```
(%i76) allMagPsi(%);
```

$$\langle I_x \rangle = 0 \quad \langle I_y \rangle = 0 \quad \langle I_z \rangle = \frac{1}{2}$$

(%o76)

7 12.3.5 Transition probabilities and the absorption of energy

The transition probability from $|\alpha\rangle$ to $|\beta\rangle$ is calculated as $\langle\beta|I_{\text{minus}}|\alpha\rangle|^2$, where I_{minus} is the lowering operator

(%i77) $\text{abs}(b_b.I_{\text{minus}}.k_a)^2;$

(%o77) 1

$|\beta\rangle$ cannot be converted to $|\beta\rangle$

(%i78) $\text{abs}(b_b.I_{\text{minus}}.k_b)^2;$

(%o78) 0

The upward transition from an arbitrary wavefunction to $|\alpha\rangle$

(%i79) $k_{\text{psi}};$

(%o79) $\begin{pmatrix} ca \\ cb \end{pmatrix}$

(%i80) $\text{abs}(b_a.I_{\text{plus}}.k_{\text{psi}})^2;$

(%o80) $|cb|^2$

Because cb is a complex number, $|cb|$ represents $cb \cdot \text{conjugate}(cb)$